

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 December 2001 (13.12.2001)

PCT

(10) International Publication Number  
**WO 01/94566 A2**

(51) International Patent Classification<sup>7</sup>: **C12N 9/00**

Woodlands, TX 77382 (US). **SANDS, Arthur, T.**; 163  
Bristol Bend Circle, The Woodlands, TX 77382 (US).

(21) International Application Number: PCT/US01/26687

(74) Agents: **ISHIMOTO, Lance, K.** et al.; LEXICON GE-  
NETICS INCORPORATED, 4000 Research Forest Drive,  
The Woodlands, TX 77381 (US).

(22) International Filing Date: 25 January 2001 (25.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/179,000 28 January 2000 (28.01.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(71) Applicant: **LEXICON GENETICS INCORPORATED**  
[US/US]; 4000 Research Forest Drive, The Woodlands, TX  
77381 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors: **DONOHU, Gregory**; 95 Autumn Branch  
Drive, The Woodlands, TX 77382 (US). **HILBUN,**  
**Erin**; 16222 Stuebner Airline, Spring, TX 77379 (US).  
**SCOVILLE, John**; 8222 Kingsbrook Dr. #543, Hous-  
ton, TX 77024 (US). **TURNER, C., Alexander, Jr.**; 67  
Winter Wheat Place, The Woodlands, TX 77381 (US).  
**FRIEDRICH, Glenn**; c/o Breland & Breland, 2207  
Hermann Drive, Houston, TX 77004 (US). **ABUIN,**  
**Alejandro**; 19 Belcarra Place, The Woodlands, TX 77382  
(US). **ZAMBROWICZ, Brian**; 18 Firethorne Place, The

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: NOVEL HUMAN ENZYMES AND POLYNUCLEOTIDES ENCODING THE SAME

(57) Abstract: Novel human polynucleotide and polypeptide sequences are disclosed that can be used in therapeutic, diagnostic, and pharmacogenomic applications.



WO 01/94566 A2

NOVEL HUMAN ENZYMES AND  
POLYNUCLEOTIDES ENCODING THE SAME

The present application claims the benefit of U.S.  
5 Provisional Application Number 60/179,000 which was filed on  
January 28, 2000 and is herein incorporated by reference in its  
entirety.

1. INTRODUCTION

10 The present invention relates to the discovery,  
identification, and characterization of novel human  
polynucleotides encoding proteins sharing sequence similarity with  
mammalian enzymes. The invention encompasses the described  
polynucleotides, host cell expression systems, the encoded  
15 protein, fusion proteins, polypeptides and peptides, antibodies to  
the encoded proteins and peptides, and sequencetically engineered  
animals that either lack or over express the disclosed sequences,  
antagonists and agonists of the proteins, and other compounds that  
modulate the expression or activity of the proteins encoded by the  
20 disclosed polynucleotides that can be used for diagnosis, drug  
screening, clinical trial monitoring and the treatment of  
physiological disorders.

2. BACKGROUND OF THE INVENTION

25 Enzymes are biological catalysts that modify biological  
substrates, including proteins, as part of degradation,  
maturation, catabolic, metabolic, differentiation, and secretory  
pathways within the body. Enzyme abnormalities have thus been  
associated with, *inter alia*, growth, development, protein and  
30 cellular senescence, cancer, or other diseases.

3. SUMMARY OF THE INVENTION

The present invention relates to the discovery,  
identification, and characterization of nucleotides that encode

novel human proteins, and the corresponding amino acid sequences of these proteins. The novel human proteins (NHPs) described for the first time herein share structural similarity with nitrilase proteins from a wide variety of living organisms.

5       The novel human nucleic acid (cDNA) sequences described herein, encode proteins/open reading frames (ORFs) of 276, 159, 121, 168, 130, 152, and 285 amino acids in length (see respectively SEQ ID NOS: 2, 4, 6, 8, 10, 12 and 14).

10       The invention also encompasses agonists and antagonists of the described NHPs including small molecules, large molecules, mutant NHPs, or portions thereof that compete with native NHPs, NHP peptides, and antibodies, as well as nucleotide sequences that can be used to inhibit the expression of the described NHPs (e.g., antisense and ribozyme molecules, and gene or regulatory sequence  
15 replacement constructs) or to enhance the expression of the described NHPs (e.g., expression constructs that place the described sequence under the control of a strong promoter system), and transgenic animals that express a NHP transgene, or "knock-outs" (which can be conditional) that do not express a functional  
20 NHP.

      Further, the present invention also relates to processes for identifying compounds that modulate, i.e., act as agonists or antagonists, of NHP expression and/or NHP activity that utilize purified preparations of the described NHP and/or NHP product, or  
25 cells expressing the same. Such compounds can be used as therapeutic agents for the treatment of any of a wide variety of symptoms associated with biological disorders or imbalances.

#### 4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

30       The Sequence Listing provides the sequences of the NHP ORFs encoding the described NHP amino acid sequences. SEQ ID NOS:15 describe representative a nitrilase-like ORF with flanking sequences.

## 5. DETAILED DESCRIPTION OF THE INVENTION

The NHPs, described for the first time herein, are novel proteins that are expressed in, *inter alia*, human cell lines, gene trapped cells and human brain, fetal brain, pituitary, cerebellum, spinal cord, thymus, spleen, lymph node, bone marrow, trachea, lung, kidney, fetal liver, liver, prostate, testis, thyroid, adrenal gland, pancreas, salivary gland, stomach, small intestine, colon, skeletal muscle, heart, placenta, mammary gland, adipose, skin, esophagus, bladder, pericardium, hypothalamus, ovary, fetal kidney, and fetal lung cells.

The described sequences were compiled from gene trapped cDNAs and clones isolated from human prostate, lymph node, pituitary, mammary gland, and kidney cDNA library (Edge Biosystems, Gaithersburg, MD). The present invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, the expression products of such nucleotides, and: (a) nucleotides that encode mammalian homologs of the described sequences, including the specifically described NHPs, and the NHP products; (b) nucleotides that encode one or more portions of a NHP that correspond to functional domains of the NHP, and the polypeptide products specified by such nucleotide sequences, including but not limited to the novel regions of any active domain(s); (c) isolated nucleotides that encode mutant versions, engineered or naturally occurring, of a described NHP in which all or a part of at least one domain is deleted or altered, and the polypeptide products specified by such nucleotide sequences, including but not limited to soluble proteins and peptides in which all or a portion of the signal sequence is deleted; (d) nucleotides that encode chimeric fusion proteins containing all or a portion of a coding region of a NHP, or one of its domains (*e.g.*, a receptor or ligand binding domain, accessory protein/self-association domain, etc.) fused to another peptide or polypeptide; or (e) therapeutic or diagnostic derivatives of the

described polynucleotides such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy constructs comprising a sequence first disclosed in the Sequence Listing.

As discussed above, the present invention includes: (a) the  
5 human DNA sequences presented in the Sequence Listing (and vectors comprising the same) and additionally contemplates any nucleotide sequence encoding a contiguous NHP open reading frame (ORF), or a contiguous exon splice junction first described in the Sequence Listing, that hybridizes to a complement of a DNA sequence  
10 presented in the Sequence Listing under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO<sub>4</sub>, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green  
15 Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at p. 2.10.3) and encodes a functionally equivalent gene product. Additionally contemplated are any nucleotide sequences that hybridize to the complement of the DNA sequence that encode and express an amino acid sequence presented in the Sequence  
20 Listing under moderately stringent conditions, e.g., washing in 0.2xSSC/0.1% SDS at 42°C (Ausubel et al., 1989, *supra*), yet still encode a functionally equivalent NHP product. Functional equivalents of a NHP include naturally occurring NHPs present in other species and mutant NHPs whether naturally occurring or  
25 engineered (by site directed mutagenesis, gene shuffling, directed evolution as described in, for example, U.S. Patent No. 5,837,458). The invention also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequences.

Additionally contemplated are polynucleotides encoding a NHP  
30 ORF, or its functional equivalent, encoded by a polynucleotide sequence that is about 99, 95, 90, or about 85 percent similar or identical to corresponding regions of the nucleotide sequences of the Sequence Listing (as measured by BLAST sequence comparison

analysis using, for example, the GCG sequence analysis package using standard default settings).

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the described NHP nucleotide sequences. Such hybridization conditions may be highly stringent or less highly stringent, as described above. In instances where the nucleic acid molecules are deoxyoligonucleotides ("DNA oligos"), such molecules are generally about 16 to about 100 bases long, or about 20 to about 80, or about 34 to about 45 bases long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence Listing. Such oligonucleotides can be used in conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc.

Alternatively, such NHP oligonucleotides can be used as hybridization probes for screening libraries, and assessing gene expression patterns (particularly using a micro array or high-throughput "chip" format). Additionally, a series of the described NHP oligonucleotide sequences, or the complements thereof, can be used to represent all or a portion of the described NHP sequences. An oligonucleotide or polynucleotide sequence first disclosed in at least a portion of one or more of the sequences of SEQ ID NOS: 1-15 can be used as a hybridization probe in conjunction with a solid support matrix/substrate (resins, beads, membranes, plastics, polymers, metal or metallized substrates, crystalline or polycrystalline substrates, etc.). Of particular note are spatially addressable arrays (i.e., gene chips, microtiter plates, etc.) of oligonucleotides and polynucleotides, or corresponding oligopeptides and polypeptides, wherein at least one of the biopolymers present on the spatially addressable array comprises an oligonucleotide or polynucleotide

sequence first disclosed in at least one of the sequences of SEQ ID NOS: 1-15, or an amino acid sequence encoded thereby. Methods for attaching biopolymers to, or synthesizing biopolymers on, solid support matrices, and conducting binding studies thereon are  
5 disclosed in, *inter alia*, U.S. Patent Nos. 5,700,637, 5,556,752, 5,744,305, 4,631,211, 5,445,934, 5,252,743, 4,713,326, 5,424,186, and 4,689,405 the disclosures of which are herein incorporated by reference in their entirety.

Addressable arrays comprising sequences first disclosed in  
10 SEQ ID NOS:1-15 can be used to identify and characterize the temporal and tissue specific expression of a gene. These addressable arrays incorporate oligonucleotide sequences of sufficient length to confer the required specificity, yet be within the limitations of the production technology. The length  
15 of these probes is within a range of between about 8 to about 2000 nucleotides. Preferably the probes consist of 60 nucleotides and more preferably 25 nucleotides from the sequences first disclosed in SEQ ID NOS:1-15.

For example, a series of the described oligonucleotide  
20 sequences, or the complements thereof, can be used in chip format to represent all or a portion of the described sequences. The oligonucleotides, typically between about 16 to about 40 (or any whole number within the stated range) nucleotides in length can partially overlap each other and/or the sequence may be  
25 represented using oligonucleotides that do not overlap. Accordingly, the described polynucleotide sequences shall typically comprise at least about two or three distinct oligonucleotide sequences of at least about 8 nucleotides in length that are each first disclosed in the described Sequence  
30 Listing. Such oligonucleotide sequences can begin at any nucleotide present within a sequence in the Sequence Listing and proceed in either a sense (5'-to-3') orientation vis-a-vis the described sequence or in an antisense orientation.

Microarray-based analysis allows the discovery of broad patterns of genetic activity, providing new understanding of gene functions and generating novel and unexpected insight into transcriptional processes and biological mechanisms. The use of addressable arrays comprising sequences first disclosed in SEQ ID NOS:1-15 provides detailed information about transcriptional changes involved in a specific pathway, potentially leading to the identification of novel components or gene functions that manifest themselves as novel phenotypes.

Probes consisting of sequences first disclosed in SEQ ID NOS:1-15 can also be used in the identification, selection and validation of novel molecular targets for drug discovery. The use of these unique sequences permits the direct confirmation of drug targets and recognition of drug dependent changes in gene expression that are modulated through pathways distinct from the drugs intended target. These unique sequences therefore also have utility in defining and monitoring both drug action and toxicity.

As an example of utility, the sequences first disclosed in SEQ ID NOS:1-15 can be utilized in microarrays or other assay formats, to screen collections of genetic material from patients who have a particular medical condition. These investigations can also be carried out using the sequences first disclosed in SEQ ID NOS:1-15 *in silico* and by comparing previously collected genetic databases and the disclosed sequences using computer software known to those in the art.

Thus the sequences first disclosed in SEQ ID NOS:1-15 can be used to identify mutations associated with a particular disease and also as a diagnostic or prognostic assay.

Although the presently described sequences have been specifically described using nucleotide sequence, it should be appreciated that each of the sequences can uniquely be described using any of a wide variety of additional structural attributes, or combinations thereof. For example, a given sequence can be



described by the net composition of the nucleotides present within a given region of the sequence in conjunction with the presence of one or more specific oligonucleotide sequence(s) first disclosed in the SEQ ID NOS: 1-15. Alternatively, a restriction map  
5 specifying the relative positions of restriction endonuclease digestion sites, or various palindromic or other specific oligonucleotide sequences can be used to structurally describe a given sequence. Such restriction maps, which are typically generated by widely available computer programs (e.g., the  
10 University of Wisconsin GCG sequence analysis package, SEQUENCHER 3.0, Gene Codes Corp., Ann Arbor, MI, etc.), can optionally be used in conjunction with one or more discrete nucleotide sequence(s) present in the sequence that can be described by the relative position of the sequence relative to one or more  
15 additional sequence(s) or one or more restriction sites present in the disclosed sequence.

For oligonucleotide probes, highly stringent conditions may refer, for example, to washing in 6xSSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for  
20 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as NHP gene antisense molecules, useful, for example, in NHP gene regulation (for and/or as antisense primers in amplification reactions of NHP nucleic acid sequences). With respect to NHP gene regulation, such techniques  
25 can be used to regulate biological functions. Further, such sequences may be used as part of ribozyme and/or triple helix sequences that are also useful for NHP gene regulation.

Inhibitory antisense or double stranded oligonucleotides can additionally comprise at least one modified base moiety which is  
30 selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine,

5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 10 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide can also comprise at least one 15 modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide will comprise at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, 20 a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an  $\alpha$ -anomeric oligonucleotide. An  $\alpha$ -anomeric oligonucleotide 25 forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual  $\beta$ -units, the strands run parallel to each other (Gautier *et al.*, 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue *et al.*, 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA 30 analogue (Inoue *et al.*, 1987, FEBS Lett. 215:327-330).

Alternatively, double stranded RNA can be used to disrupt the expression and function of a targeted NHP.

Oligonucleotides of the invention can be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate  
5 oligonucleotides can be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

10 Low stringency conditions are well known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual (and  
15 periodic updates thereof), Cold Springs Harbor Press, N.Y.; and Ausubel et al., 1989, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y.

Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using appropriately  
20 stringent conditions or by PCR. The identification and characterization of human genomic clones is helpful for identifying polymorphisms (including, but not limited to, nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single nucleotide polymorphisms),  
25 determining the genomic structure of a given locus/allele, and designing diagnostic tests. For example, sequences derived from regions adjacent to the intron/exon boundaries of the human gene can be used to design primers for use in amplification assays to detect mutations within the exons, introns, splice sites (e.g.,  
30 splice acceptor and/or donor sites), etc., that can be used in diagnostics and pharmacogenomics.

Further, a NHP homolog can be isolated from nucleic acid from an organism of interest by performing PCR using two degenerate or

"wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be total RNA, mRNA, and/or cDNA obtained by reverse transcription of mRNA prepared from human or  
5 non-human cell lines or tissue known or suspected to express an allele of a NHP gene.

The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequence of the desired NHP sequence. The PCR fragment can then be used to isolate a full  
10 length cDNA clone by a variety of methods. For example, the amplified fragment can be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment can be used to isolate genomic clones via the screening of a genomic library.

15 PCR technology can also be used to isolate full length cDNA sequences. For example, RNA can be isolated, following standard procedures, from an appropriate cellular or tissue source (i.e., one known, or suspected, to express a NHP sequence, such as, for example, testis tissue). A reverse transcription (RT) reaction  
20 can be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second  
25 strand synthesis may then be primed with a complementary primer. Thus, cDNA sequences upstream of the amplified fragment can be isolated. For a review of cloning strategies that can be used, see e.g., Sambrook et al., 1989, *supra*.

A cDNA encoding a mutant NHP sequence can be isolated, for  
30 example, by using PCR. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in an individual putatively carrying a mutant NHP allele, and by

extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR, 5 optionally cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant NHP allele to that of a corresponding normal NHP allele, the mutation(s) responsible for the loss or alteration of function of the mutant 10 NHP gene product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known to carry a mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, obesity, high blood pressure, 15 connective tissue disorders, infertility, etc.), or a cDNA library can be constructed using RNA from a tissue known, or suspected, to express a mutant NHP allele. A normal NHP sequence, or any suitable fragment thereof, can then be labeled and used as a probe to identify the corresponding mutant NHP allele in such libraries. 20 Clones containing mutant NHP sequences can then be purified and subjected to sequence analysis according to methods well known to those skilled in the art.

Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a 25 tissue known, or suspected, to express a mutant NHP allele in an individual suspected of or known to carry such a mutant allele. In this manner, gene products made by the putatively mutant tissue can be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against normal 30 NHP product, as described below. (For screening techniques, see, for example, Harlow, E. and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.)

Additionally, screening can be accomplished by screening with labeled NHP fusion proteins, such as, for example, alkaline phosphatase-NHP or NHP-alkaline phosphatase fusion proteins.

In cases where a NHP mutation results in an expressed gene product with altered function (e.g., as a result of a missense or a frameshift mutation), polyclonal antibodies to NHP are likely to cross-react with a corresponding mutant NHP gene product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well known in the art.

The invention also encompasses (a) DNA vectors that contain any of the foregoing NHP coding sequences and/or their complements (i.e., antisense); (b) DNA expression vectors that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences (for example, baculo virus as described in U.S. Patent No. 5,869,336 herein incorporated by reference); (c) genetically engineered host cells that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (d) genetically engineered host cells that express an endogenous NHP sequence under the control of an exogenously introduced regulatory element (i.e., gene activation). As used herein, regulatory elements include, but are not limited to, inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include but are not limited to the cytomegalovirus (hCMV) immediate early gene, regulatable, viral elements (particularly retroviral LTR promoters), the early or late promoters of SV40 adenovirus, the *lac* system, the *trp* system, the *TAC* system, the *TRC* system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate

kinase (PGK), the promoters of acid phosphatase, and the promoters of the yeast  $\alpha$ -mating factors.

The present invention also encompasses antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists and agonists of a NHP, as well as compounds or nucleotide constructs that inhibit expression of a NHP gene (transcription factor inhibitors, antisense and ribozyme molecules, or gene or regulatory sequence replacement constructs), or promote the expression of a NHP (e.g., expression constructs in which NHP coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.).

The NHPs or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, antibodies, antagonists and agonists can be useful for the detection of mutant NHPs or inappropriately expressed NHPs for the diagnosis of disease. The NHP proteins or peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for screening for drugs (or high throughput screening of combinatorial libraries) effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of a NHP in the body. The use of engineered host cells and/or animals may offer an advantage in that such systems allow not only for the identification of compounds that bind to the endogenous receptor for a NHP, but can also identify compounds that trigger NHP-mediated activities or pathways.

Finally, the NHP products can be used as therapeutics. For example, soluble derivatives such as NHP peptides/domains corresponding to NHP, NHP fusion protein products (especially NHP-Ig fusion proteins, i.e., fusions of a NHP, or a domain of a NHP, to an IgFc), NHP antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate or act on downstream targets in a NHP-

mediated pathway) can be used to directly treat diseases or disorders. For instance, the administration of an effective amount of soluble NHP, or a NHP-IgFc fusion protein or an anti-idiotypic antibody (or its Fab) that mimics the NHP could activate  
5 or effectively antagonize the endogenous NHP receptor. Nucleotide constructs encoding such NHP products can be used to genetically engineer host cells to express such products *in vivo*; these genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of a NHP, a NHP peptide, or a NHP  
10 fusion protein to the body. Nucleotide constructs encoding functional NHP, mutant NHPs, as well as antisense and ribozyme molecules can also be used in "gene therapy" approaches for the modulation of NHP expression. Thus, the invention also encompasses pharmaceutical formulations and methods for treating  
15 biological disorders.

Various aspects of the invention are described in greater detail in the subsections below.

#### 5.1 THE NHP SEQUENCES

20 The cDNA sequences and the corresponding deduced amino acid sequences of the described NHP are presented in the Sequence Listing. SEQ ID NOS:15 describes the NHP ORFs as well as flanking regions. The NHP nucleotides were obtained from human cDNA libraries using probes and/or primers generated from human gene  
25 trapped sequence tags. Expression analysis has provided evidence that the described NHP can be expressed in a variety of human cells as well as gene trapped human cells.

#### 5.2 NHP AND NHP POLYPEPTIDES

30 NHPs, polypeptides, peptide fragments, mutated, truncated, or deleted forms of the NHPs, and/or NHP fusion proteins can be prepared for a variety of uses. These uses include, but are not limited to, the generation of antibodies, as reagents in



diagnostic assays, for the identification of other cellular gene products related to a NHP, as reagents in assays for screening for compounds that can be as pharmaceutical reagents useful in the therapeutic treatment of mental, biological, or medical disorders and disease.

The Sequence Listing discloses the amino acid sequence encoded by the described NHP polynucleotides. The NHPs display initiator methionines in DNA sequence contexts consistent with translation initiation sites, and apparently display signal sequences which can indicate that the described NHP ORFs are secreted proteins or possibly membrane associated.

The NHP amino acid sequences of the invention include the amino acid sequences presented in the Sequence Listing as well as analogues and derivatives thereof, as well as any oligopeptide sequence of at least about 10-40, generally about 12-35, or about 16-30 amino acids in length first disclosed in the Sequence Listing. Further, corresponding NHP homologues from other species are encompassed by the invention. In fact, any NHP encoded by the NHP nucleotide sequences described above are within the scope of the invention, as are any novel polynucleotide sequences encoding all or any novel portion of an amino acid sequence presented in the Sequence Listing. The degenerate nature of the genetic code is well known, and, accordingly, each amino acid presented in the Sequence Listing, is generically representative of the well known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such, as contemplated herein, the amino acid sequences presented in the Sequence Listing, when taken together with the genetic code (see, for example, Table 4-1 at page 109 of "Molecular Cell Biology", 1986, J. Darnell et al. eds., Scientific American Books, New York, NY, herein incorporated by reference) are generically representative of all the various permutations and combinations of nucleic acid sequences that can encode such amino acid sequences.

The invention also encompasses proteins that are functionally equivalent to the NHPs encoded by the presently described nucleotide sequences as judged by any of a number of criteria, including, but not limited to, the ability to bind and cleave a substrate of a NHP, or the ability to effect an identical or complementary downstream pathway, or a change in cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation, etc.). Such functionally equivalent NHP proteins include, but are not limited to, additions or substitutions of amino acid residues within the amino acid sequence encoded by the NHP nucleotide sequences described above, but which result in a silent change, thus producing a functionally equivalent gene product. Amino acid substitutions can be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the invention. Where, as in the present instance, the NHP products or NHP polypeptides can be produced in soluble or secreted forms (by removing one or more transmembrane domains where applicable), the peptide or polypeptide can be recovered from the culture media. Such expression systems also encompass engineered host cells that express a NHP, or a functional equivalent, *in situ*. Purification or enrichment of NHP from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well known to those skilled in the art. However, such

engineered host cells themselves may be used in situations where it is important not only to retain the structural and functional characteristics of the NHP, but to assess biological activity, e.g., in drug screening assays.

5       The expression systems that may be used for purposes of the invention include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing NHP nucleotide sequences; yeast (e.g., *Saccharomyces*,  
10 *Pichia*) transformed with recombinant yeast expression vectors containing NHP encoding nucleotide sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing NHP sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower  
15 mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing NHP nucleotide sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression  
20 constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

      In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the  
25 NHP product being expressed. For example, when a large quantity of such a protein is to be produced for the generation of pharmaceutical compositions of or containing NHP, or for raising antibodies to a NHP, vectors that direct the expression of high levels of fusion protein products that are readily purified may be  
30 desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which a NHP coding sequence may be ligated

individually into the vector in frame with the *lacZ* coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, *Nucleic Acids Res.* 13:3101-3109; Van Heeke & Schuster, 1989, *J. Biol. Chem.* 264:5503-5509); and the like. pGEX  
5 vectors (Pharmacia or American Type Culture Collection) can also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the  
10 presence of free glutathione. The PGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

In an insect system, *Autographa californica* nuclear polyhydrosis virus (AcNPV) is used as a vector to express foreign  
15 sequences. The virus grows in *Spodoptera frugiperda* cells. A NHP coding sequence can be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of NHP coding sequence will  
20 result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (*i.e.*, virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted sequence is expressed (*e.g.*, see  
25 Smith et al., 1983, *J. Virol.* 46: 584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the NHP nucleotide sequence of interest may  
30 be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric sequence may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-

essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing a NHP product in infected hosts (e.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific  
5 initiation signals may also be required for efficient translation of inserted NHP nucleotide sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire NHP sequence or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate  
10 expression vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the  
15 reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer  
20 elements, transcription terminators, etc. (See Bittner et al., 1987, Methods in Enzymol. 153:516-544).

In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such  
25 modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell  
30 lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript,

glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, human cell lines.

5 For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the NHP sequences described above can be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with  
10 DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are  
15 switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer  
20 cell lines which express the NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the NHP product.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et  
25 *al.*, 1977, *Cell* 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, *Proc. Natl. Acad. Sci. USA* 48:2026), and adenine phosphoribosyltransferase (Lowy, *et al.*, 1980, *Cell* 22:817) genes can be employed in *tk*<sup>-</sup>, *hgp*<sup>-</sup> or *aprt*<sup>-</sup> cells, respectively. Also,  
30 antimetabolite resistance can be used as the basis of selection for the following genes: *dhfr*, which confers resistance to methotrexate (Wigler, *et al.*, 1980, *Natl. Acad. Sci. USA* 77:3567; O'Hare, *et al.*, 1981, *Proc. Natl. Acad. Sci. USA* 78:1527); *gpt*,

which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150:1); and hygro, which confers resistance to  
5 hygromycin (Santerre, et al., 1984, Gene 30:147).

Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins  
10 expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-8976). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts  
15 from cells infected with recombinant vaccinia virus are loaded onto Ni<sup>2+</sup>-nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

Also encompassed by the present invention are fusion proteins that direct the NHP to a target organ and/or facilitate transport  
20 across the membrane into the cytosol. Conjugation of NHPs to antibody molecules or their Fab fragments could be used to target cells bearing a particular epitope. Attaching the appropriate signal sequence to the NHP would also transport the NHP to the desired location within the cell. Alternatively targeting of NHP  
25 or its nucleic acid sequence might be achieved using liposome or lipid complex based delivery systems. Such technologies are described in Liposomes: A Practical Approach, New, RRC ed., Oxford University Press, New York and in U.S. Patents Nos. 4,594,595, 5,459,127, 5,948,767 and 6,110,490 and their respective  
30 disclosures which are herein incorporated by reference in their entirety. Additionally embodied are novel protein constructs engineered in such a way that they facilitate transport of the NHP to the target site or desired organ, where they cross the cell

membrane and/or the nucleus where the NHP can exert its functional activity. This goal may be achieved by coupling of the NHP to a cytokine or other ligand that provides targeting specificity, and/or to a protein transducing domain (see generally U.S.

5 applications Ser. No. 60/111,701 and 60/056,713, both of which are herein incorporated by reference, for examples of such transducing sequences) to facilitate passage across cellular membranes and can optionally be engineered to include nuclear localization sequences.

### 10 5.3 ANTIBODIES TO NHP PRODUCTS

Antibodies that specifically recognize one or more epitopes of a NHP, or epitopes of conserved variants of a NHP, or peptide fragments of a NHP are also encompassed by the invention. Such antibodies include but are not limited to polyclonal antibodies, 15 monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')<sub>2</sub> fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

20 The antibodies of the invention may be used, for example, in the detection of NHP in a biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of NHP. Such antibodies may also be utilized in conjunction with, for example, 25 compound screening schemes for the evaluation of the effect of test compounds on expression and/or activity of a NHP gene product. Additionally, such antibodies can be used in conjunction gene therapy to, for example, evaluate the normal and/or engineered NHP-expressing cells prior to their introduction into 30 the patient. Such antibodies may additionally be used as a method for the inhibition of abnormal NHP activity. Thus, such antibodies may, therefore, be utilized as part of treatment methods.



For the production of antibodies, various host animals may be immunized by injection with the NHP, an NHP peptide (e.g., one corresponding to a functional domain of an NHP), truncated NHP polypeptides (NHP in which one or more domains have been deleted),  
5 functional equivalents of the NHP or mutated variant of the NHP. Such host animals may include but are not limited to pigs, rabbits, mice, goats, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to  
10 Freund's adjuvant (complete and incomplete), mineral salts such as aluminum hydroxide or aluminum phosphate, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*.  
15 Alternatively, the immune response could be enhanced by combination and or coupling with molecules such as keyhole limpet hemocyanin, tetanus toxoid, diphtheria toxoid, ovalbumin, cholera toxin or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the  
20 sera of the immunized animals.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, can be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not  
25 limited to, the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal  
30 Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated *in vitro* or

*in vivo*. Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison *et al.*, 1984, *Proc. Natl. Acad. Sci.*, 81:6851-6855; Neuberger *et al.*, 1984, *Nature*, 312:604-608; Takeda *et al.*, 1985, *Nature*, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. Such technologies are described in U.S. Patents Nos. 6,075,181 and 5,877,397 and their respective disclosures which are herein incorporated by reference in their entirety. Also favored is the production of fully humanized monoclonal antibodies as described in US Patent No. 6,150,584 and respective disclosures which are herein incorporated by reference in their entirety.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778; Bird, 1988, *Science* 242:423-426; Huston *et al.*, 1988, *Proc. Natl. Acad. Sci. USA* 85:5879-5883; and Ward *et al.*, 1989, *Nature* 334:544-546) can be adapted to produce single chain antibodies against NHP gene products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, such fragments include, but are not limited to: the F(ab')<sub>2</sub> fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed (Huse *et al.*, 1989, *Science*,

246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Antibodies to a NHP can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" a given NHP, using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, 1993, FASEB J 7(5):437-444; and Nissinoff, 1991, J. Immunol. 147(8):2429-2438). For example antibodies which bind to a NHP domain and competitively inhibit the binding of NHP to its cognate receptor can be used to generate anti-idiotypes that "mimic" the NHP and, therefore, bind and activate or neutralize a receptor. Such anti-idiotypic antibodies or Fab fragments of such anti-idiotypes can be used in therapeutic regimens involving a NHP signaling pathway.

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims. All cited publications, patents, and patent applications are herein incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising at least 24 contiguous bases of nucleotide sequence first disclosed  
5 in SEQ ID NO: 1.
2. An isolated nucleic acid molecule comprising a nucleotide sequence that:
  - (a) encodes the amino acid sequence shown in SEQ ID  
10 NO: 2; and
  - (b) hybridizes under stringent conditions to the nucleotide sequence of SEQ ID NO: 1 or the complement thereof.
- 15 3. An isolated nucleic acid molecule encoding the amino acid sequence described in SEQ ID NO: 2.
4. An isolated oligopeptide comprising at least about 12 amino acids in a sequence first disclosed in SEQ ID NO:2.  
20

## SEQUENCE LISTING

&lt;110&gt; LEXICON GENETICS INCORPORATED

<120> Novel Human Enzymes and Polynucleotides  
Encoding the Same

&lt;130&gt; LEX-0118-PCT

&lt;150&gt; US 60/179,000

&lt;151&gt; 2000-01-28

&lt;160&gt; 15

&lt;170&gt; FastSEQ for Windows Version 4.0

&lt;210&gt; 1

&lt;211&gt; 831

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 1

```

atgacctctt tccgcttggc cctcatccag cttcagattt cttccatcaa atcagataac      60
gtcactcgcg cttgtagctt catccgggag gcagcaacgc aaggagccaa aatagtttct      120
ttgccggaat gctttaattc tccatatgga gcgaaatatt ttcctgaata tgcagagaaa      180
attcctgggtg aatccacaca gaagctttct gaagtagcaa aggaatgcag catatatctc      240
attggaggct ctatccctga agaggatgct gggaaattat ataacacctg tgctgtgttt      300
gggcctgatg gaactttact agcaaagtat agaaagatcc atctgtttga cattgatgtt      360
cctggaaaaa ttacatttca agaattctaaa acattgagtc cgggtgatag tttctccaca      420
tttgatactc cttactgcag agtgggtctg ggcactctgct acgacatgcg gtttgcagag      480
cttgcacaaa tctacgcaca gagaggctgc cagctgttgg tatatccagg agcttttaat      540
ctgaccactg gaccagccca ttgggagtta cttcagcgaa gccgggctgt tgataatcag      600
gtgtatgtgg ccacagctc tcctgcccg gtagcaaaag cctctatgt tgctgggga      660
cacagcaccg tgggtgaacc ttggggggag gttctagcca aagctggcac agaagaagca      720
atcgtgtatt cagacataga cctgaagaag ctggctgaaa tacgccagca aatccccgtt      780
tttagacaga agcgatcaga cctctatgct gtggagatga aaaagcccta a      831

```

&lt;210&gt; 2

&lt;211&gt; 276

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

```

Met Thr Ser Phe Arg Leu Ala Leu Ile Gln Leu Gln Ile Ser Ser Ile
  1              5              10              15
Lys Ser Asp Asn Val Thr Arg Ala Cys Ser Phe Ile Arg Glu Ala Ala
      20              25              30
Thr Gln Gly Ala Lys Ile Val Ser Leu Pro Glu Cys Phe Asn Ser Pro
      35              40              45
Tyr Gly Ala Lys Tyr Phe Pro Glu Tyr Ala Glu Lys Ile Pro Gly Glu
      50              55              60
Ser Thr Gln Lys Leu Ser Glu Val Ala Lys Glu Cys Ser Ile Tyr Leu
      65              70              75              80
Ile Gly Gly Ser Ile Pro Glu Glu Asp Ala Gly Lys Leu Tyr Asn Thr
      85              90              95

```

Cys Ala Val Phe Gly Pro Asp Gly Thr Leu Leu Ala Lys Tyr Arg Lys  
 100 105 110  
 Ile His Leu Phe Asp Ile Asp Val Pro Gly Lys Ile Thr Phe Gln Glu  
 115 120 125  
 Ser Lys Thr Leu Ser Pro Gly Asp Ser Phe Ser Thr Phe Asp Thr Pro  
 130 135 140  
 Tyr Cys Arg Val Gly Leu Gly Ile Cys Tyr Asp Met Arg Phe Ala Glu  
 145 150 155 160  
 Leu Ala Gln Ile Tyr Ala Gln Arg Gly Cys Gln Leu Leu Val Tyr Pro  
 165 170 175  
 Gly Ala Phe Asn Leu Thr Thr Gly Pro Ala His Trp Glu Leu Leu Gln  
 180 185 190  
 Arg Ser Arg Ala Val Asp Asn Gln Val Tyr Val Ala Thr Ala Ser Pro  
 195 200 205  
 Ala Arg Asp Asp Lys Ala Ser Tyr Val Ala Trp Gly His Ser Thr Val  
 210 215 220  
 Val Asn Pro Trp Gly Glu Val Leu Ala Lys Ala Gly Thr Glu Glu Ala  
 225 230 235 240  
 Ile Val Tyr Ser Asp Ile Asp Leu Lys Lys Leu Ala Glu Ile Arg Gln  
 245 250 255  
 Gln Ile Pro Val Phe Arg Gln Lys Arg Ser Asp Leu Tyr Ala Val Glu  
 260 265 270  
 Met Lys Lys Pro  
 275

<210> 3  
 <211> 480  
 <212> DNA  
 <213> Homo sapiens

<400> 3  
 atgtcatgga ggatttcccc tgccacacca tgctgtaggg agttaacttt tcatttgtgc 60  
 attttctgtt tggaaacagc ttactgcaga gtgggtctgg gcatctgcta cgacatgcgg 120  
 ttgacagagc ttgcacaaat ctacgcacag agaggtctgcc agctgttggt atatccagga 180  
 gcttttaatc tgaccactgg accagcccat tgggagttac ttcagcgaag cggggtgtt 240  
 gataatcagg tgtatgtggc cacagcctct cctgcccggt atgacaaagc ctctatgtt 300  
 gcctggggac acagcacctt ggtgaaccct tggggggagg ttctagccaa agctggcaca 360  
 gaagaagcaa tcgtgtattc agacatagac ctgaagaagc tggctgaaat acgccagcaa 420  
 atccccgttt ttagacagaa gcgatcagac ctctatgctg tggagatgaa aaagccctaa 480

<210> 4  
 <211> 159  
 <212> PRT  
 <213> Homo sapiens

<400> 4  
 Met Ser Trp Arg Ile Ser Pro Ala Thr Pro Cys Cys Arg Glu Leu Thr  
 1 5 10 15  
 Phe His Leu Cys Ile Phe Cys Leu Glu Thr Ala Tyr Cys Arg Val Gly  
 20 25 30  
 Leu Gly Ile Cys Tyr Asp Met Arg Phe Ala Glu Leu Ala Gln Ile Tyr  
 35 40 45  
 Ala Gln Arg Gly Cys Gln Leu Leu Val Tyr Pro Gly Ala Phe Asn Leu  
 50 55 60  
 Thr Thr Gly Pro Ala His Trp Glu Leu Leu Gln Arg Ser Arg Ala Val  
 65 70 75 80

Asp Asn Gln Val Tyr Val Ala Thr Ala Ser Pro Ala Arg Asp Asp Lys  
                   85                  90                  95  
 Ala Ser Tyr Val Ala Trp Gly His Ser Thr Val Val Asn Pro Trp Gly  
                   100                  105                  110  
 Glu Val Leu Ala Lys Ala Gly Thr Glu Glu Ala Ile Val Tyr Ser Asp  
                   115                  120                  125  
 Ile Asp Leu Lys Lys Leu Ala Glu Ile Arg Gln Gln Ile Pro Val Phe  
                   130                  135                  140  
 Arg Gln Lys Arg Ser Asp Leu Tyr Ala Val Glu Met Lys Lys Pro  
                   145                  150                  155

<210> 5  
 <211> 366  
 <212> DNA  
 <213> Homo sapiens

<400> 5  
 atgcggtttg cagagcttgc acaaactctac gcacagagag gctgccagct gttggtatat 60  
 ccaggagctt ttaatctgac cactggacca gccattggg agttacttca gcgaagccgg 120  
 gctgttgata atcagggtga tgtggccaca gcctctcctg ccggggatga caaagcctcc 180  
 tatgttgctt ggggacacag caccgtggtg aacccttggg gggaggttct agccaaagct 240  
 ggcacagaag aagcaatcgt gtattcagac atagacctga agaagctggc tgaaatacgc 300  
 cagcaaatcc ccgttttttag acagaagcga tcagacctct atgctgtgga gatgaaaaag 360  
 ccctaa 366

<210> 6  
 <211> 121  
 <212> PRT  
 <213> Homo sapiens

<400> 6  
 Met Arg Phe Ala Glu Leu Ala Gln Ile Tyr Ala Gln Arg Gly Cys Gln  
   1                  5                  10                  15  
 Leu Leu Val Tyr Pro Gly Ala Phe Asn Leu Thr Thr Gly Pro Ala His  
                   20                  25                  30  
 Trp Glu Leu Leu Gln Arg Ser Arg Ala Val Asp Asn Gln Val Tyr Val  
                   35                  40                  45  
 Ala Thr Ala Ser Pro Ala Arg Asp Asp Lys Ala Ser Tyr Val Ala Trp  
                   50                  55                  60  
 Gly His Ser Thr Val Val Asn Pro Trp Gly Glu Val Leu Ala Lys Ala  
                   65                  70                  75                  80  
 Gly Thr Glu Glu Ala Ile Val Tyr Ser Asp Ile Asp Leu Lys Lys Leu  
                   85                  90                  95  
 Ala Glu Ile Arg Gln Gln Ile Pro Val Phe Arg Gln Lys Arg Ser Asp  
                   100                  105                  110  
 Leu Tyr Ala Val Glu Met Lys Lys Pro  
                   115                  120

<210> 7  
 <211> 507  
 <212> DNA  
 <213> Homo sapiens

<400> 7  
 atgtcatgga ggatttcccc tgccacacca tgctgtaggg agttaacttt tcatttgtgc 60  
 attttctggt tgqaaacagc ttactgcaga gtgggtctgg gcatctgcta cgacatgcgg 120

```

tttgcagagc ttgcacaaat ctacgcacag agaggctgcc agctgttggt atatccagga 180
gctttttaatc tgaccactgg accagcccat tgggagttac ttcagcgaag ccgggctggt 240
gataatcagg tgtatgtggc cacagcctct cctgcccggg atgacaaagc ctcctatggt 300
gcctggggac acagcacctg ggtgaaccct tggggggagg ttctagccaa agctggcaca 360
gaagaagcaa tcgtgtattc agacatagac ctgaagaagc tggctgaaat acgccagcaa 420
atccccgttt ttagacagaa gcgaaatatt ttcctgaata tgcagagaaa attcctgggt 480
aatccacaca gaagctttct gaagtag 507

```

&lt;210&gt; 8

&lt;211&gt; 168

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 8

```

Met Ser Trp Arg Ile Ser Pro Ala Thr Pro Cys Cys Arg Glu Leu Thr
  1             5             10             15
Phe His Leu Cys Ile Phe Cys Leu Glu Thr Ala Tyr Cys Arg Val Gly
      20             25             30
Leu Gly Ile Cys Tyr Asp Met Arg Phe Ala Glu Leu Ala Gln Ile Tyr
      35             40             45
Ala Gln Arg Gly Cys Gln Leu Leu Val Tyr Pro Gly Ala Phe Asn Leu
      50             55             60
Thr Thr Gly Pro Ala His Trp Glu Leu Leu Gln Arg Ser Arg Ala Val
      65             70             75             80
Asp Asn Gln Val Tyr Val Ala Thr Ala Ser Pro Ala Arg Asp Asp Lys
      85             90             95
Ala Ser Tyr Val Ala Trp Gly His Ser Thr Val Val Asn Pro Trp Gly
      100            105            110
Glu Val Leu Ala Lys Ala Gly Thr Glu Glu Ala Ile Val Tyr Ser Asp
      115            120            125
Ile Asp Leu Lys Lys Leu Ala Glu Ile Arg Gln Gln Ile Pro Val Phe
      130            135            140
Arg Gln Lys Arg Asn Ile Phe Leu Asn Met Gln Arg Lys Phe Leu Val
      145            150            155            160
Asn Pro His Arg Ser Phe Leu Lys
      165

```

&lt;210&gt; 9

&lt;211&gt; 393

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 9

```

atgcggtttg cagagcttgc acaaattctac gcacagagag gctgccagct gttggtatat 60
ccaggagctt ttaatctgac cactggacca gccattggg agttacttca gcgaagccgg 120
gctgttgata atcagggtga tgtggccaca gcctctcctg cccgggatga caaagcctcc 180
tatgttgctt ggggacacag caccgtgggt aacccttggg gggaggttct agccaaagct 240
ggcacagaag aagcaatcgt gtattcagac atagacctga agaagctggc tgaaatacgc 300
cagcaaatcc ccgttttttag acagaagcga aatattttcc tgaatatgca gagaaaattc 360
ctggtgaatc cacacagaag ctttctgaag tag 393

```

&lt;210&gt; 10

&lt;211&gt; 130

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens



&lt;400&gt; 10

```

Met Arg Phe Ala Glu Leu Ala Gln Ile Tyr Ala Gln Arg Gly Cys Gln
 1           5           10           15
Leu Leu Val Tyr Pro Gly Ala Phe Asn Leu Thr Thr Gly Pro Ala His
      20           25           30
Trp Glu Leu Leu Gln Arg Ser Arg Ala Val Asp Asn Gln Val Tyr Val
      35           40           45
Ala Thr Ala Ser Pro Ala Arg Asp Asp Lys Ala Ser Tyr Val Ala Trp
      50           55           60
Gly His Ser Thr Val Val Asn Pro Trp Gly Glu Val Leu Ala Lys Ala
      65           70           75           80
Gly Thr Glu Glu Ala Ile Val Tyr Ser Asp Ile Asp Leu Lys Lys Leu
      85           90           95
Ala Glu Ile Arg Gln Gln Ile Pro Val Phe Arg Gln Lys Arg Asn Ile
      100          105          110
Phe Leu Asn Met Gln Arg Lys Phe Leu Val Asn Pro His Arg Ser Phe
      115          120          125
Leu Lys
      130

```

&lt;210&gt; 11

&lt;211&gt; 459

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 11

```

atgacctctt tccgcttggc cctcatccag cttcagattt cttccatcaa atcagataac      60
gtcactcgcg cttgtagctt catccgggag gcagcaacgc aaggagccaa aatagtttct      120
ttgccggaat gctttaattc tccatatgga gcgaaatatt ttctgaata tgcagagaaa      180
attcctgggtg aatccacaca gaagctttct gaagtagcaa aggaatgcag catatatctc      240
attggaggct ctatccctga agaggatgct gggaaattat ataacacctg tgctgtgttt      300
gggcctgatg gaactttact agcaaagtat agaaagatcc atctgtttga cattgatgtt      360
cctggaaaaa ttacatttca agaattctaaa acattgagtc cgggtgatag tttctccaca      420
tttgatactc gtatgtacca gataagtttg cctcttttag      459

```

&lt;210&gt; 12

&lt;211&gt; 152

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 12

```

Met Thr Ser Phe Arg Leu Ala Leu Ile Gln Leu Gln Ile Ser Ser Ile
 1           5           10           15
Lys Ser Asp Asn Val Thr Arg Ala Cys Ser Phe Ile Arg Glu Ala Ala
      20           25           30
Thr Gln Gly Ala Lys Ile Val Ser Leu Pro Glu Cys Phe Asn Ser Pro
      35           40           45
Tyr Gly Ala Lys Tyr Phe Pro Glu Tyr Ala Glu Lys Ile Pro Gly Glu
      50           55           60
Ser Thr Gln Lys Leu Ser Glu Val Ala Lys Glu Cys Ser Ile Tyr Leu
      65           70           75           80
Ile Gly Gly Ser Ile Pro Glu Glu Asp Ala Gly Lys Leu Tyr Asn Thr
      85           90           95
Cys Ala Val Phe Gly Pro Asp Gly Thr Leu Leu Ala Lys Tyr Arg Lys
      100          105          110
Ile His Leu Phe Asp Ile Asp Val Pro Gly Lys Ile Thr Phe Gln Glu

```

115                      120                      125  
 Ser Lys Thr Leu Ser Pro Gly Asp Ser Phe Ser Thr Phe Asp Thr Arg  
 130                      135                      140  
 Met Tyr Gln Ile Ser Leu Pro Leu  
 145                      150

<210> 13  
 <211> 858  
 <212> DNA  
 <213> Homo sapiens

<400> 13  
 atgacctctt tccgcttggc cctcatccag cttcagattt cttccatcaa atcagataac 60  
 gtcactcgcg cttgtagctt catccgggag gcagcaacgc aaggagccaa aatagtttct 120  
 ttgccggaat gctttaattc tccatatgga gcgaaatatt ttcctgaata tgcagagaaa 180  
 attcctgggtg aatccacaca gaagctttct gaagtagcaa aggaatgcag catatatctc 240  
 attggaggct ctatccctga agaggatgct gggaaattat ataacacctg tgctgtgttt 300  
 gggcctgatg gaactttact agcaaagtat agaaagatcc atctgtttga cattgatgtt 360  
 cctggaaaaa ttacatttca agaatactaaa acattgagtc cgggtgatag tttctccaca 420  
 tttgatactc cttactgcag agtgggtctg ggcattctgct acgacatgcg gtttgcagag 480  
 cttgcacaaa tctacgcaca gagaggctgc cagctgttgg tatatccagg agcttttaat 540  
 ctgaccactg gaccagccca ttgggagtta cttcagcgaa gccgggctgt tgataatcag 600  
 gtgtatgtgg ccacagcctc tctgccccgg gatgacaaag cctcctatgt tgcctgggga 660  
 cacagcaccg tgggtgaacc ttggggggag gttctagcca aagctggcac agaagaagca 720  
 atcgtgtatt cagacataga cctgaagaag ctggctgaaa tacgccagca aatccccgtt 780  
 tttagacaga agcgaaatat tttcctgaat atgcagagaa aattcctggt gaatccacac 840  
 agaagctttc tgaagtag 858

<210> 14  
 <211> 285  
 <212> PRT  
 <213> Homo sapiens

<400> 14  
 Met Thr Ser Phe Arg Leu Ala Leu Ile Gln Leu Gln Ile Ser Ser Ile  
 1                      5                      10                      15  
 Lys Ser Asp Asn Val Thr Arg Ala Cys Ser Phe Ile Arg Glu Ala Ala  
 20                      25                      30  
 Thr Gln Gly Ala Lys Ile Val Ser Leu Pro Glu Cys Phe Asn Ser Pro  
 35                      40                      45  
 Tyr Gly Ala Lys Tyr Phe Pro Glu Tyr Ala Glu Lys Ile Pro Gly Glu  
 50                      55                      60  
 Ser Thr Gln Lys Leu Ser Glu Val Ala Lys Glu Cys Ser Ile Tyr Leu  
 65                      70                      75                      80  
 Ile Gly Gly Ser Ile Pro Glu Glu Asp Ala Gly Lys Leu Tyr Asn Thr  
 85                      90                      95  
 Cys Ala Val Phe Gly Pro Asp Gly Thr Leu Leu Ala Lys Tyr Arg Lys  
 100                      105                      110  
 Ile His Leu Phe Asp Ile Asp Val Pro Gly Lys Ile Thr Phe Gln Glu  
 115                      120                      125  
 Ser Lys Thr Leu Ser Pro Gly Asp Ser Phe Ser Thr Phe Asp Thr Pro  
 130                      135                      140  
 Tyr Cys Arg Val Gly Leu Gly Ile Cys Tyr Asp Met Arg Phe Ala Glu  
 145                      150                      155                      160  
 Leu Ala Gln Ile Tyr Ala Gln Arg Gly Cys Gln Leu Leu Val Tyr Pro  
 165                      170                      175

Gly Ala Phe Asn Leu Thr Thr Gly Pro Ala His Trp Glu Leu Leu Gln  
 180 185 190  
 Arg Ser Arg Ala Val Asp Asn Gln Val Tyr Val Ala Thr Ala Ser Pro  
 195 200 205  
 Ala Arg Asp Asp Lys Ala Ser Tyr Val Ala Trp Gly His Ser Thr Val  
 210 215 220  
 Val Asn Pro Trp Gly Glu Val Leu Ala Lys Ala Gly Thr Glu Glu Ala  
 225 230 235 240  
 Ile Val Tyr Ser Asp Ile Asp Leu Lys Lys Leu Ala Glu Ile Arg Gln  
 245 250 255  
 Gln Ile Pro Val Phe Arg Gln Lys Arg Asn Ile Phe Leu Asn Met Gln  
 260 265 270  
 Arg Lys Phe Leu Val Asn Pro His Arg Ser Phe Leu Lys  
 275 280 285

<210> 15  
 <211> 3093  
 <212> DNA  
 <213> Homo sapiens

<400> 15  
 ggatggtggg gcatacctgt ggtcccagct acataagagg ctgagacaag aggattgcct 60  
 gaactgagta ggtcaaggct gcagtggacc atgtttgtgc cactgcactc cagcctgggc 120  
 gacagaacaa ggccctgcct caaaataaaa aatattagct aatggaaagt gattatcata 180  
 aaagctaaaa gggaacttta aagaacagaa gaaaagcaaa tatgatgtat agctactacc 240  
 tccaggaaga aataagcttg gaagagcccc caacctcctt gctccagggc tgagcacaga 300  
 ccttgctcagg gctggctaca taatttgtgg ggcccagttc ccttgttcag atagcaagag 360  
 aaaagtgtcg ttagcttttc cttctgcagt atctctttca acctctcatg gtgttatttg 420  
 ctgtttaatg tcatgttctc ttggacacat gaatacttat ggggtaagtg cagactttta 480  
 gaggtgcctg ggaccctgt cctgtgaata ggcatgtgtg cagctcactg gctgccaggt 540  
 ttccctctg ccagcagcgg gatcgatgtg ctgtgaccca gccagtagtg gggaaactga 600  
 gacagacatc ttcccttccc atgagctggg cctgctcatg ggaattatgt gaggcagctc 660  
 caaggaatca cactttctgt gctgggacat actcaagtat atggattgga ggtagacgag 720  
 agggccattg aacaaacagt aagggacagg accatattca aaccagtcct ttttacttta 780  
 agccatattc ctcatctcat tcccctacac tgcgtagtaa gaagctggtt cactctagat 840  
 tcttgctcct ggcattgggac tttgcccatt gatattgtc tatctccaga tagattttag 900  
 actattgaca ttttgacag gataattctt cgttggtgta tggagggggg tgtcctatgc 960  
 attgtaggat gtttggcagt atccttggtc tctattcatt agatgccact catacctcat 1020  
 cagttgtggc atcaaaggta tcttcagaca ttgtcagatg tccccccggg gacataactg 1080  
 ccttccattt gagaactatg gctctgtctg aatccagcag ttcgatcttc tgatagctgt 1140  
 tttcttttgt ctttgttctc agcccccccc cccccggtag gacccgcggg ccgcccggatc 1200  
 tccagcgtc agtccgcgcc gcagggtgtg cttgtctgca gagtcatgac ctctttccgc 1260  
 ttggccctca tccagcttca gatttcttcc atcaaatcag ataacgtcac tcgcgcttgt 1320  
 agcttcatcc gggaggcagc aacgcaagga gccaaaatag tttctttgcc ggaatgcttt 1380  
 aattctccat atggagcgaa atattttcct gaatatgcag agaaaattcc tggatgaatcc 1440  
 acacagaagc tttctgaagt agcaaaggaa tgcagcatat atctcattgg aggctctatc 1500  
 cctgaagagg atgctgggaa attatataac acctgtgctg tgtttgggcc tgatggaact 1560  
 ttactagcaa agtatagaaa gatccatctg ttgacattg atgttcctgg aaaaattaca 1620  
 ttcaagaat ctaaaacatt gagtccgggt gatagtttct ccacatttga tactcgtatg 1680  
 taccagataa gtttgccctt ttagcaatct cagtagaaga caatcaggta tttatttctt 1740  
 ttttgtctct ctccgatttc ttcacataac ctaactgaaa gaccataagt gagaaaggca 1800  
 gagaatcatc acagatctgg aaagttcggg cttattttgag aactaaggat ttgacacgat 1860  
 tttgcccctt gatttgattg tagcttctctg ttacggcttc cagagtatac ctattaggct 1920  
 acagttgagt acctcccatc tagataataa gcattcaatt agaataaatt tctcatcttt 1980  
 actccgctga tgtaaatgat gtctttatga gatgaagtcc aagtaggaat gagcttgtaa 2040  
 attatctctg tctcaggtc ctgtgttaat ttatccctgt cagtgttttg tgatcattat 2100

gtcatggagg	atttcccctg	ccacaccatg	ctgtagggag	ttaacttttc	atttgtgcat	2160
tttctgtttg	gaaacagctt	actgcagagt	gggtctgggc	atctgctacg	acatgcggtt	2220
tgcagagctt	gcacaaatct	acgcacagag	aggctgccag	ctgttggtat	atccaggagc	2280
ttttaatctg	accactggac	cagcccattg	ggagttactt	cagcgaagcc	gggctgttga	2340
taatcaggtg	tatgtggcca	cagcctctcc	tgccgggat	gacaaagcct	cctatgttgc	2400
ctggggacac	agcaccgtgg	tgaacccttg	gggggaggtt	ctagccaaag	ctggcacaga	2460
agaagcaatc	gtgtattcag	acatagacct	gaagaagctg	gctgaaatac	gccagcaaat	2520
ccccgttttt	agacagaagc	gatcagacct	ctatgctgtg	gagatgaaaa	agccctaaag	2580
tttatgtttc	taatgtgtca	cagaatagga	cgatatgatt	ctacaacata	atcaactccc	2640
tattaaattc	tttaatgaag	aaaaaaaaaa	aaaaaaaaaa	aaaaaatatt	ttcctgaata	2700
tgcagagaaa	attcctggtg	aatccacaca	gaagctttct	gaagtagcaa	aggaatgcag	2760
catatatctc	attggaggct	ctatccctga	agaggatgct	gggaaattat	ataacacctg	2820
tgctgtgttt	gggcctgatg	gaactttact	agcaaagtat	agaaagatcc	atctgtttga	2880
cattgatgtt	cctggaaaaa	ttacatttca	agaatctaaa	acattgagtc	cgggtgatag	2940
tttctccaca	tttgatactc	cttactgcag	agtgggtctg	ggcatctgct	acgacatgcg	3000
gtttgcagag	cttgcacaaa	tctacgcaca	gagaggctgc	cagctgttgg	tatatccagg	3060
agcttttaat	ctgaccactg	gaccagccca	ttg			3093